arXiv:1209.1604v1 [cond-mat.quant-gas] 7 Sep 2012

Comment on "Hydrodynamic fluctuations and the minimum shear viscosity of the dilute Fermi gas at unitarity"

Paul Romatschke and Ryan Edward Young

Department of Physics, 390 UCB, University of Colorado, Boulder, CO 80309, USA

Abstract

We confirm and expand on work by Chafin and Schäfer on hydrodynamic fluctuations in the unitary Fermi gas. Using the result for the equation of state from an recent MIT experiment, we derive lower bounds for η/n and η/s as a function of temperature. Re-analyzing recent quantum Monte Carlo data for the shear-viscosity spectral function we point out a possible resolution for the tension between the viscosity bound $\eta/n \gtrsim 0.3$ from Chafin and Schäfer and the quantum Monte Carlo results $\eta/n \lesssim 0.2$ from Wlazlowski et al. near the critical temperature.

I. MOTIVATION

Hydrodynamics allows for long lived shear and sound waves. Fluctuations can populate these modes and interaction of these waves can dissipate momentum. This contributes to the shear viscosity and other transport coefficients.

Our study builds upon results found in the context of relativistic fluids, specifically Ref. [1] which was recently applied to the case of the Fermi gas at unitarity by Chafin and Schäfer in Ref. [2]. In Ref. [2] it was found that the dimensionless combination shear viscosity over density η/n must be larger or equal to 0.32 for the unitary Fermi gas at the superfluid-normal phase transition temperature $T = T_c$. Thus, there is a tension between this result and the result for η/n found in the quantum Monte Carlo simulations by Wlazlowski et al. in Ref. [3], who find $\eta/n|_{T\sim T_c} \lesssim 0.2$. This provides the motivation for us to re-derive and generalize the result by Chafin and Schäfer as well as re-analyze the results by Wlazlowski et al.

II. DERIVATION OF THE CORRELATOR

We will be interested in a particular retarded correlator of the energy-stress tensor $T^{\mu\nu}$, namely

$$G_R^{xyxy}(x-y) = \langle T^{xy}(x) T^{xy}(y)\rangle_R\,,$$

because the Fourier transform of this correlator in *classical hydrodynamics* (meaning in the absence of fluctuations) can be related to the value of transport coefficients. Specifically, for the case of a conformal system with vanishing bulk viscosity coefficient, one has ([2], cf. [4])

$$G_R^{xyxy}(\omega,\mathbf{k}) = P - i\eta\omega + \eta\tau_R\omega^2 - \frac{\kappa}{2}\mathbf{k}^2 + \mathcal{O}(\omega^3,\mathbf{k}^3)\,, \tag{1}$$

with η the shear viscosity coefficient and τ_R the relaxation time. It is possible to derive this result within the stochastic formulation of hydrodynamics (see e.g. Ref. [5] for a recent review). The key idea is to supplement the energy-stress tensor with a noise term $\xi^{\mu\nu}$ that is modeling fluctuations,

$$T^{\mu\nu} \to T^{\mu\nu}_{\rm cl} + \xi^{\mu\nu}\,, \quad \langle \xi^{\mu\nu}(x)\xi^{\alpha\beta}(y)\rangle \propto \delta^4(x-y)\,,$$

where the index "cl" refers to purely classical hydrodynamic expressions and $\langle\rangle$ here denotes the noise average (see appendix A for an illustration). In the case of the G^{xyxy} correlator, it has been shown in Ref. [1] that to leading order in fluctuations, the stochastic formulation reproduces the $-i\omega\eta$ term in Eq. (1), while to next-to-leading (one-loop) order, the result can be written as

$$
\begin{aligned}
\left[G_R^{xyxy}\right]^{1-loop}(\omega, k=0) \; = \; & \rho^2 \int \frac{d\omega'}{2\pi} \int \frac{d^3\mathbf{p}}{(2\pi)^3} \times \\
& \Big(\Delta_S^{xx}(\omega', \mathbf{p}) \Delta_R^{yy}(\omega-\omega', -\mathbf{p}) + \Delta_S^{xy}(\omega', \mathbf{p}) \Delta_R^{yx}(\omega-\omega', -\mathbf{p}) + \\
& \Delta_S^{xx}(\omega', \mathbf{p}) \Delta_R^{yy}(\omega-\omega', -\mathbf{p}) + \Delta_S^{xy}(\omega', \mathbf{p}) \Delta_R^{yx}(\omega-\omega', -\mathbf{p}) \Big) \,, \qquad (2)
\end{aligned}
$$

where $\Delta_{S,R}^{ij}$ are the symmetric and retarded version of the velocity-velocity correlator (see appendix A). In the zero momentum limit $|\mathbf{k}| \to 0$, the integrations in the one-loop contribution to the correlator are straightforward (cf. [1]) and one finds

$$
\left[G_R^{xyxy}\right]^{1-loop}(\omega \ll p_{\rm max}, |\mathbf{k}|=0) = -i\omega \frac{17 T p_{max}}{120\gamma\pi^2} + (1+i)\omega^{3/2} \frac{(28+3\sqrt{6})T}{980\pi\gamma^{3/2}} + \ldots \,, \qquad (3)
$$

where $\gamma = \frac{\eta}{\rho}$ and $p_{\rm max}$ is the cut-off for the effective theory of hydrodynamics. Note that terms not included in Eq. (3) are those that are suppressed by higher powers of $p_{\rm max}\gamma$, which is the "small parameter" of the effective theory. Similarly, higher loop contributions to G_R^{xyxy} are suppressed by higher powers of $p_{\rm max}\gamma$. Thus, up to first order in powers of $p_{\rm max}\gamma$, the result for the retarded correlator is given by the sum of the "tree-level" result Eq. (1) and the one-loop result Eq. (3).

III. MINIMUM ON VISCOSITY

Using the Kubo relation

$$
\eta = \lim_{\omega \to 0} \frac{-{\rm Im} G_R^{xyxy}(\omega, \mathbf{k}=0)}{\omega} \,,
$$

the result for the physical viscosity including fluctuations to first loop order is given by

$$
\eta = \eta_{\rm cl} + \frac{17 T p_{\rm max} \rho}{120 \eta_{\rm cl} \pi^2} \,,
$$

where again the index "cl" refers to "classical" contributions in the sense of turning off all fluctuations. As was pointed out, our calculation is systematic if

$$
p_{\rm max} \ll \frac{\rho}{\eta_{\rm cl}} \,,
$$

so we parametrize $p_{\max}$ by

$$p_{\max} = \frac{\rho}{\eta_{\rm cl}}\Phi\,,$$

where Φ is a dimensionless function of thermodynamic variables that should be small. For instance, we may take[1] Φ to be a function of pressure over density, $\Phi = \Phi(P/\rho)$. Then the result for the physical viscosity may be cast into a result for the dimensionless viscosity over density ratio,

$$\left(\frac{\eta}{n}\right) = \left(\frac{\eta}{n}\right)_{\rm cl} + \frac{17T\rho^2\Phi(P/\rho)}{120n^3\left(\eta/n\right)_{\rm cl}^2\pi^2}\,.$$

This expression can be extremized, finding that the physical viscosity over density expression may never decrease below the minimum value

$$\left(\frac{\eta}{n}\right) \geq \left(\frac{\eta}{n}\right)_{\rm min} = \left(\frac{153T\rho^2\Phi(P/\rho)}{160\pi^2n^3}\right)^{1/3}\,. \tag{4}$$

The minimum value for viscosity is dimensionless and presumably universal, meaning that for a unitary Fermi gas it should only depend on the combination T/T_F with $T_F = \frac{(3\pi^2 n)^{2/3}}{2m}$. Assuming a power-like behavior, we have

$$\left(\frac{153T\rho^2\Phi(P/\rho)}{160\pi^2n^3}\right) \propto \left(\frac{T}{T_F}\right)^{\alpha}\,,$$

with α a constant. Similarly, we take Φ to be given by a power law behavior with

$$\Phi(P/\rho) \propto \left(P/\rho\right)^{\beta}\,,$$

with β a constant. As shown in appendix B, the pressure is given by the relation $P = nTG(T/T_F)$ and $\lim_{T\to\infty} G = 1$. With this, we find

$$\frac{Tm^2}{n}\left(\frac{T}{m}\right)^{\beta} \propto \left(\frac{T}{T_F}\right)^{\alpha}\,,$$

which leads to the requirement $\alpha = 1+\beta\,,\ 2-\beta = \alpha$ and hence

$$\alpha = \frac{3}{2}\,, \quad \beta = \frac{1}{2}\,.$$

Thus we obtain for the final form of the lower viscosity bound

$$\left(\frac{\eta}{n}\right) \geq \left(\frac{\eta}{n}\right)_{\rm min} = \left(k\frac{153T\rho^2\sqrt{P/\rho)}}{160\pi^2n^3}\right)^{1/3}\,, \tag{5}$$

---

[1] In principle, Φ may also depend on the combination T/T_F with T_F the Fermi temperature (see below). However, we will drop this dependence in favor of a constant that may vary around unity.

with k an arbitrary constant of order unity. One finds

$$\left(\frac{\eta}{n}\right) \geq \left(\frac{\eta}{n}\right)_{\rm min} = \left(k\frac{459\sqrt{\frac{T^3}{T_F^3}\frac{P}{nT}}}{2^{3/2}160}\right)^{1/3} \simeq 1.005k^{1/3}\sqrt{\frac{T}{T_F}}\left(\frac{P}{nT}\right)^{1/6}\,, \tag{6}$$

which matches the result found in Ref. [2].

IV. WIDTH OF SPECTRAL FUNCTION PEAK

Collecting all terms of the retarded correlator $G_R^{xyxy}(\omega)$ from Eq.(1) and Eq. (3), the full result for the one-loop spectral function $\eta(\omega)$ to leading order in frequency is given by

$$\begin{aligned}\frac{\eta(\omega)}{n} &= \left(\frac{\eta}{n}\right)_{\rm cl} + k\frac{51}{2^{3/2}120}\left(\frac{\eta}{n}\right)_{\rm cl}^{-2}\left(\frac{T}{T_F}\right)^{3/2}\sqrt{\frac{P}{nT}}\\ &\quad -\frac{3\pi(28+3\sqrt{6})}{2^{3/2}980}\left(\frac{\eta}{n}\right)_{\rm cl}^{-3/2}\left(\frac{T}{T_F}\right)\sqrt{\frac{\omega}{T_F}}+\mathcal{O}(\omega^2)\,,\end{aligned} \tag{7}$$

with k an unknown constant of order unity. Let us now assume we have arrived at the viscosity coefficient $\lim_{\omega\to 0}\frac{\eta(\omega)}{n}$ by some other means, e.g. a first principle lattice calculation.

The viscosity coefficient then fixes the value of $\eta_{\rm cl}/n$ and hence Eq. (7) gives the low-frequency behavior of the spectral function. We may then compare this result to the spectral function obtained by the first-principles method.

Let perform this analysis on the spectral function for the shear viscosity obtained with the Path Integral Monte Carlo (PIMC) approach from Ref. [3]. In this article, the authors analyzed data using the Maximum Entropy Method (among others) based on the 5-parameter model class

$$\eta(\hat\omega, m, \sigma, c, \alpha_1, \alpha_2) = f(\hat\omega, \alpha_1, \alpha_2)\frac{C}{15\pi\sqrt{\hat\omega}} + \left(1 - f(\hat\omega, \alpha_1, \alpha_2)\right)N(\hat\omega, m, \sigma, c)\,,$$

where $f = e^{-\alpha_1\alpha_2}\frac{e^{\alpha_1\hat\omega}-1}{1+e^{\alpha_1(\hat\omega-\alpha_s)}}$, $N = \frac{c}{\sqrt{2\pi\sigma^2}}e^{-(\hat\omega-m)^2/2/\sigma^2}$, $\hat\omega = \omega/T_F$ and C is Tan's contact (which is determined by other means and hence will be assumed to be known). This class of models implements the known large frequency behavior because $f(\omega\to\infty)\to 1$. Interestingly, note that the model also is compatible with the low frequency behavior in Eq. (7) because $f(\hat\omega \ll 1) = \frac{\alpha_1}{1+e^{\alpha_1\alpha_2}}\hat\omega$. It is therefore possible that this model reproduces the correct low-frequency behavior, even though it is not guaranteed. We re-analyze the data from

Ref. [3] for the correlator $G(\tau)$ for the $N_x = 8, N_\tau = 52$ simulation on $T/T_F = 0.355$ using a simple χ^2 minimization of

$$G(\tau) - \frac{1}{\pi}\int_0^\infty d\hat{\omega}\eta(\hat{\omega}, m, \sigma, c, \alpha_1, \alpha_2)\hat{\omega}\frac{\cosh(\hat{\omega}(\tau T_F - \frac{T_F}{2T}))}{\sinh(\hat{\omega}T_F/2/T)} \to 0\,, \tag{8}$$

neglecting the first three values of τ for reasons explained in Ref. [3]. We find a very small $\chi^2/\mathrm{dof} \sim 4 \times 10^{-3}$, but also observe that the minimum is very broad in parameter space (χ^2 does not change appreciably when changing parameters by order one). The extracted spectral function is shown in Fig. 1(a). As can be seen from this figure, in our re-analysis of the PIMC data, the extracted shape of $\eta(\omega)$ is close to the result reported in Ref. [3], but the tail asymptotics are only matched at much higher values of frequency. Nevertheless, the low frequency behavior, including the extracted viscosity coefficient $\eta \equiv \lim_{\omega\to 0}\eta(\omega)$ is rather similar. Using the viscosity coefficient to extract a value for quantity η_{cl}/n as described above (assuming $k \simeq 1$), we can then compare the PIMC results to the purely hydrodynamic result Eq. (7)[2]. We find that for the case of $T/T_F = 0.355$, the spectral function shape extracted from the PIMC data agrees reasonably well with the behavior demanded from Eq. (7). Because it matches the known low frequency behavior, we conclude that this provides additional evidence that the extracted value of the shear viscosity of a unitary Fermi gas at $T/T_F = 0.35$ is $\eta/n \simeq 0.7$.

Let us now repeat our analysis for the low temperature case $T/T_F = 0.15$. Simple minimization of Eq. (8) leads to the result shown in Fig. 1(b). Comparison to the result from Ref. [3] now indicates that both analysis agree with each other for large frequencies, while there is considerable discrepancy at low frequencies. While Ref. [3] finds a spectral function that is essentially featureless at low frequency, our re-analysis of the same data show a peak at $\omega = 0$. Trying to convert the viscosity coefficient $\eta(\omega = 0)$ into a value for η_{cl}/n we find that in both cases, the constant k has to be less than unity. In particular, for our re-analysis we need $k \simeq 0.5$, while to reach the viscosity value found in Ref. [3], $k \sim 0.05$ is needed. The latter value is a much larger deviation from unity than could be expected. Nevertheless, with these choices for k, we can evaluate the low frequency behavior of the spectral function, finding the results shown in Fig. 1(b). For our re-analysis,

---

[2] In general, the extraction of η_{cl}/n will not be single-valued. While negative values of η_{cl}/n are clearly disallowed, we choose the smallest non-negative value of η_{cl}/n.



FIG. 1: Spectral function $\eta(\omega)/n$ extracted from PIMC simulation Ref. [3] with $N_x = 8$, $N_\tau = 52$ for $T/T_F = 0.355$ (plot (a)) and $T/T_F = 0.15$ (plot (b)). Dots: (approximate) result published in Ref. [3]. Dashed: our re-analysis of the same data using a simple χ^2 fit (Eq. (8)). Also shown is the known UV behavior ("high ω asymptotics"). Solid lines are low frequency behavior extracted from the known hydrodynamic behavior Eq. (7). Since the $T/T_F = 0.15$ low frequency peak is extremely narrow, we have highlighted it using an ellipse in plot (b).

the low frequency shape is roughly consistent with the hydrodynamic behavior, although the agreement is somewhat less than satisfactory.

For the results for the viscosity coefficient from Ref. [3], we find that the hydrodynamic behavior in Eq. (7) results in an extremely narrow peak, in complete disagreement with the spectral function shape found in Ref. [3]. We suspect that if indeed such a narrow peak is present in the physical spectral function, it could have been missed by the method used in Ref. [3] to reconstruct the spectral function from the imaginary time correlation function. If we simply add on such a low frequency peak to the broad spectral function found in Ref. [3], we find that the resulting value for the shear viscosity at $T/T_F = 0.15$ is $\eta/n \simeq 0.35$, consistent with our re-analysis of the PIMC data, and consistent with the viscosity bound found in Ref. [2].



FIG. 2: Summary of results for η/n (left) and η/s (right). The viscosity limit from Eq. (6) was used to delineate a “disallowed region” of viscosity values using the MIT equation of state from Ref. [9] for $P/n/T$ and S/N. Also shown are results from kinetic theory at high temperatures and the phonon scattering at low temperatures (see text for details), as well as results from Enss et al., Ref. [8]. Furthermore, we indicate the results from the PIMC calculation by Wlazlowski et al. [3] and our own re-analysis of the same data. Note that the resulting value for η/n is consistent with the “upper bound” reported in the supplement to Ref. [3].

V. DISCUSSION AND CONCLUSION

Employing the framework of hydrodynamics in the presence of fluctuations, the low frequency behaviour to the shear viscosity spectral function has been calculated. The calculation gives a bound on the physical shear viscosity, Eq. (6), which matches the result found in Ref. [2] but disagrees with the low-temperature results from Ref. [3]. We pointed out that the reason for the discrepancy can be tracked to the low-frequency shape of the spectral function extracted in Ref. [3]. Specifically, assuming that the shape from Ref. [3] is correct for $\omega > 0.1T_F$, hydrodynamic fluctuations predict a narrow but high peak at $\omega < 0.1T_F$, which effectively doubles the viscosity value. The situation is summarized in Fig. 2, where the viscosity bound (6) is shown together with the viscosity in kinetic theory (cf. Ref [6])

$$\left.\frac{\eta}{n}\right|_{KT} = \frac{45\pi^{3/2}}{32\times 2^{3/2}}\left(\frac{T}{T_F}\right)^{3/2} \simeq 2.768\left(\frac{T}{T_F}\right)^{3/2} \tag{9}$$

as well as the viscosity of phonon scattering in the superfluid phase, (cf. Ref [7])

$$\left.\frac{\eta}{n}\right|_{phonons} = 5.3 \times 10^{-5} \xi^{7/2} \left(\frac{T_F}{T}\right)^5 \simeq 1.76 \times 10^{-6} \left(\frac{T_F}{T}\right)^5 . \qquad (10)$$

We conclude that once the low frequency peak in the spectral function is included in the analysis, the viscosity value from the PIMC method no longer is in conflict with the lower viscosity limit (6). Also, note that in this case the viscosity value are then much closer to the result from Enss et al. reported in Ref. [8].

Also shown in Fig. 2 are results for the minimum η/s, using the experimental data from Ref. [9] for the entropy per particle as a function of temperature. The corresponding results for kinetic theory and phonons are [11],[10]

$$\left.\frac{\eta}{s}\right|_{KT} = 2.768 \left(\frac{T}{T_F}\right)^{3/2} \frac{1}{\frac{5}{2} + \ln\left[\frac{3\sqrt{\pi}}{4}(T/T_F)^{3/2}\right]} , \quad \left.\frac{\eta}{s}\right|_{phonons} = 5.94 \times 10^{-8} \left(\frac{T_F}{T}\right)^8 . \qquad (11)$$

One observes that the viscosity result from kinetic theory enters the disallowed region at around $T/T_F \lesssim 0.3$. We interpret this as meaning that kinetic theory looses its validity at or above this temperature. Similarly, the viscosity result from the superfluid analysis enters the disallowed region for $T/T_F \gtrsim 0.1$. Again, we interpret this as meaning that the calculation looses validity at or below this temperature. The results for the shear viscosity by Enss et al. stay within the allowed region for all temperatures shown, but it is interesting to note that the η/s values at low temperatures stop at the boundary to the disallowed region.

We conclude that the theory of hydrodynamic fluctuations can provide a relevant tool to study the properties of unitary Fermi gases. Specifically, we expect that our findings may help in the analysis of future quantum Monte Carlo calculations. We plan to further develop this theory in the future.

Acknowledgments

We would like to thank T. Schäfer for sharing an early version of Ref. [2] with us, as well as for fruitful discussion. Furthermore, we would like to thank M. Zwierlein and M. Ku for providing us with the experimental data from Ref. [9]. Moreover, we would like to thank G. Wlazlowski for providing us the PIMC data for the correlator from Ref. [3] and pointing out the supplemental material included in Ref. [3]. Finally, we would like to thank T. Enss for providing us with the results from Ref. [8].

Appendix A: Finding the Velocity-Velocity Correlator

Let us start by writing down the Navier-Stokes and continuity equations

$$\frac{\partial}{\partial t}(\rho v_k) + \frac{\partial \Pi_{ik}}{\partial x_k} = 0\,, \quad \frac{\partial}{\partial t}\rho + \frac{\partial}{\partial x_k}(\rho v_k) = 0\,, \tag{A1}$$

where Π_{ik} is the standard viscous fluid stress tensor. For consistency with conformal field theories the bulk viscosity, ξ, has been set to zero. This is a good approximation of cold atom systems. Ultimately, we would like to quantify how thermal fluctuations dissipate momentum, acting to increase viscosity. This requires finding the velocity-velocity correlator, so that the retarded correlator of two stress tensors can be found. In order to find the velocity-velocity correlator, a stochastic formulation of fluid dynamics is needed. To do this a noise term, Ξ_{ij}, is added to the stress tensor, $\Pi_{ik} \to \Pi_{ik} + \Xi_{ij}$. The choice of noise term will dictate what form of fluid dynamics is found. To reproduce the Navier-Stokes equations, the noise must have a short distance Gaussian correlation.

$$\Xi^{jk} = \frac{1}{2}(\xi^{jk} + \xi^{kj} - \frac{2}{3}\delta_{jk}\xi^{ll}) \tag{A2}$$

$$\langle \xi^{ik}\xi^{nm}\rangle = \eta T(\delta_{il}\delta_{km} + \delta_{kl}\delta_{im})\delta(x - x') \tag{A3}$$

Now with this stochastic formulation we can find the hydrodynamic correlation functions. Consider a system in equilibrium, allow for small perturbations away from equilibrium in v and $\rho \to \rho_o + \delta\rho$. The equation of fluid mechanics to first order in the perturbations are

$$\frac{\partial}{\partial t}(\rho v_k) + \frac{\partial}{\partial x_k}(\Pi_{ik} + \Xi_{ik}) = 0\,, \quad \partial_t \rho + \rho_0 \frac{\partial v^k}{\partial x^k} = 0\,. \tag{A4}$$

The velocity can be decomposed into longitudinal and transverse parts satisfying

$$\nabla \cdot v_t(r,t) = 0\,, \quad \nabla \times v_l(r,t) = 0\,. \tag{A5}$$

These restrictions give a new system of fluid dynamic equations. The linear form of the continuity equation yields

$$\frac{\partial \rho}{\partial t} + \rho_0 \nabla \cdot v_l = 0 \tag{A6}$$

and momentum conservation yields the following system of equations

$$\frac{\partial}{\partial t} v_t = \frac{\eta}{\rho_0}\Delta v_t\,, \quad \rho_0 \frac{\partial}{\partial t} v_l = -\nabla \delta P + \frac{4}{3}\eta \nabla(\nabla \cdot v_l)\,. \tag{A7}$$

Assuming that the velocity $v(r,t)$ goes rapidly to zero at large distances, we can define the Laplace-Fourier transform

$$v(k,\omega) = \int dr \int_0^\infty d\omega \; e^{-ik\cdot r + i\omega t} v(r,t) \tag{A8}$$

In an infinite medium the Laplace-Fourier transform allows us to solve the equations of motion. Considering only the transverse velocity

$$v_t = \frac{k\xi_{ij}}{-i\omega + \gamma k^2} \tag{A9}$$

one has

$$\langle v_i^t v_j^t \rangle = (\delta_{ij} - \frac{k_i k_j}{k^2}) \frac{k^2 \langle (\xi_{ij})^2 \rangle}{\omega^2 + \gamma^2 k^4} \tag{A10}$$

with $\gamma = \eta/\rho$, the kinematic viscosity. It follows that

$$\langle v_i^t v_j^t \rangle = \frac{2T}{\rho} (\delta_{ij} - \frac{k_i k_j}{k^2}) \frac{\gamma k^2}{\omega^2 + \gamma^2 k^4} \,. \tag{A11}$$

Repeating this process for the longitudinal velocity, the entire velocity-velocity correlator is found to be

$$\langle v_i v_j \rangle = \frac{2T}{\rho} \Big[\frac{k_i k_j}{k^2} \frac{\tilde{\gamma}\omega^2 k^2}{(\omega^2 - c^2 k^2)^2 + (\tilde{\gamma}\omega^2 k^2)^2} + \Big(\delta_{ij} - \frac{k_i k_j}{k^2} \Big) \frac{\gamma k^2}{\omega^2 + \gamma^2 k^4} \Big] \equiv \Delta_S^{ij} \tag{A12}$$

where $\tilde{\gamma} = \frac{4}{3}\gamma$ and we have introduced the symbol Δ_S for the symmetric velocity-velocity correlator. Now let us calculate the retarded velocity-velocity correlator. The classical Fluctuation-dissipation theorem reads

$$\chi''(\omega) = \frac{1}{2} \beta \omega C(\omega) \tag{A13}$$

with $\beta = 1/T$, being the inverse temperature and

$$C_{ij}(t) = \langle A_i(t) A_j(0) \rangle_c = \langle \delta A_i(t) \delta A_j(0) \rangle \,. \tag{A14}$$

This is the Kubo or relaxation function with the subscript c stands for cumulant or connected. In Fourier space $C_{ij}(\omega)$ is the symmetric correlator of the fluctuations.

To find the retarded correlator, we must examine the generalized susceptibility. It is related to the time derivative of the Kubo function by

$$\chi_{ij}(t) = -\beta \Theta(t) \dot{C}_{ij}(t) \tag{A15}$$



FIG. 3: Left: results for the normalized pressure P/P_0 from the MIT experiment Ref. [9] and a 5 parameter fit (see text for details). Right: The function $G = \frac{P}{nT}$ as a function of T/T_F.

where $\Theta(t)$ is a step-function function. The generalized susceptibility χ can be identified as the retarded correlator. Utilizing the relation $\chi'' = \mathrm{Im}(\chi)$ and that χ is analytic in the upper half plane we can find the retarded velocity correlation function:

$$\Delta_R^{ij} = \frac{1}{\rho}\Big[\frac{k_i k_j}{k^2}\frac{\omega^2}{(\omega^2 - c^2k^2)^2 + (i\omega k^2\tilde{\gamma})} + \Big(\delta_{ij} - \frac{k_i k_j}{k^2}\Big)\frac{-\gamma k^2}{-i\omega + \gamma k^2}\Big] \tag{A16}$$

The Kubo-Martin-Schwinger conditions relate the symmetric and retarded correlator in quantum mechanical regimes. In Fermi systems the response functions are related to bosonic operator (e.g. bilinears of fermi operators). Therefore, a classical version of the KMS relation is found to be $G_s = \frac{2T}{\omega}\mathrm{Im}(G_R)$.

Appendix B: Equation of State

The equation of state of a cold unitary Fermi gas was measured experimentally by the MIT group [9]. The result from Ref. [9] for the pressure (versus fugacity $\zeta = e^{-\mu/T}$) is shown in Fig. 3(a). The result for the pressure is normalized by the ideal non-interacting single-component Fermi gas pressure,

$$P_0(\mu, T) = -T\lambda_{dB}^{-3}\mathrm{Li}_{5/2}(-\zeta^{-1})\,,$$

where $\lambda_{dB} = \sqrt{\frac{2\pi}{mT}}$ is the deBroglie wavelength. Also shown in Fig. 3(a) is a fit of the experimental data using the parametrization

$$h(\zeta) \equiv P(\mu,T)/P_0(\mu,T) = \frac{\zeta^3 + c_1\zeta^2 + c_2\zeta + c_3}{\zeta^3 + c_4\zeta^2 + c_5\zeta + 1}$$

which is a variant of the approach followed in Ref. [12]. A least square fit gives

$$c_1 = 101.591\,, \quad c_2 = 158.375\,, \quad c_3 = 104.884\,, \quad c_4 = 4.30128\,, \quad c_5 = 67.3894\,.$$

Note that c_4 is related to the Bertsch parameter ξ as $c_4 = \xi^{-3/2}$ from the relation $P(\mu, T = 0) = \xi^{-3/2} P_0(\mu, T = 0)$. The fitted value of c_4 therefore corresponds to $\xi = 0.378$. Using the thermodynamic relation $n = \frac{\partial P}{\partial \mu}\Big|_T$ and the identity $n = \lambda_{dB}^{-3}\frac{(4\pi)^{3/2}}{3\pi^2}\left(\frac{T_F}{T}\right)^{3/2}$ one has

$$\frac{(4\pi)^{3/2}}{3\pi^2}\left(\frac{T_F}{T}\right)^{3/2} = \zeta\frac{\partial}{\partial\zeta}\left(\mathrm{Li}_{5/2}(-\zeta^{-1})h(\zeta)\right)\,,$$

and hence a (numerical) result for $\zeta = \zeta(T/T_F)$ by inverting the above functional relation.

Using the above relations the pressure may be written as

$$P(\mu,T) = nTG(\zeta)\,, \quad G(\zeta) = \frac{1}{-\zeta\frac{d}{d\zeta}\ln\left[-\mathrm{Li}_{5/2}(-\zeta^{-1})h(\zeta)\right]}\,.$$

For convenience, the function G is plotted in Fig. 3(b) as a function of T/T_F. As can be seen, the form of G implies that the pressure is close to nT, but not monotonically larger or smaller than nT in the domain considered. We have checked that $\lim_{T\to\infty} G(T/T_F) \to 1$, as expected.

[1] P. Kovtun, G. D. Moore and P. Romatschke, Phys. Rev. D **84** (2011) 025006 [arXiv:1104.1586 [hep-ph]].

[2] C. Chafin and T. Schaefer, arXiv:1209.1006 [cond-mat.quant-gas].

[3] G. Wlazlowski, P. Magierski and J. E. Drut, Phys. Rev. Lett. **109** (2012) 020406 [arXiv:1204.0270 [cond-mat.quant-gas]].

[4] R. Baier, P. Romatschke, D. T. Son, A. O. Starinets and M. A. Stephanov, JHEP **0804** (2008) 100 [arXiv:0712.2451 [hep-th]].

[5] P. Kovtun, arXiv:1205.5040 [hep-th].

[6] P. Massignan, G. M. Bruun and H. Smith, Phys. Rev. A **71** (2005) 033607 [cond-mat/0409660 [cond-mat.stat-mech]].

[7] M. Braby, J. Chao and T. Schafer, Phys. Rev. A **82** (2010) 033619 [arXiv:1003.2601 [cond-mat.quant-gas]].

[8] T. Enss, R. Haussmann and W. Zwerger, Annals Phys. **326** (2011) 770 [arXiv:1008.0007 [cond-mat.quant-gas]].

[9] M.J.H. Ku, A.T. Sommer, L.W. Cheuk and M.W. Zwierlein, Science, **335** (2012) 563. [arXiv:1110.3309 [cond-mat.quant-gas]].

[10] G. Rupak and T. Schafer, Phys. Rev. A **76** (2007) 053607 [arXiv:0707.1520 [cond-mat.other]].

[11] T. Schafer and D. Teaney, Rept. Prog. Phys. **72** (2009) 126001 [arXiv:0904.3107 [hep-ph]].

[12] T. Schafer, Phys. Rev. A **82** (2010) 063629 [arXiv:1008.3876 [cond-mat.quant-gas]].